

*Review***The role of the melatoninergetic system in epilepsy and comorbid psychiatric disorders**

Jana Tchekalarova^{a*}, Slavianka Moyanova^b, Antonio De Fusco^b, Richard Teke Ngomba^b

^a Institute of Neurobiology, Acad. G. Bonchev Str., Bl. 23, Bulgarian Academy of Sciences,
Sofia 1113, Bulgaria

^b I.R.C.C.S. NEUROMED, Località Camerelle, Pozzilli 86077 (IS), Italy

*Corresponding author: Jana Tchekalarova, Institute of Neurobiology, Acad. G. Bonchev Str.,
Bl. 23, Bulgarian Academy of Sciences, Sofia 1113, Bulgaria.

Contents

Abstract.....	00
1. Introduction.....	00
2. The melatonin system and anxiety.....	00
3. The melatonin system and depression.....	00
4. The melatonin system and epilepsy.....	00
4.1. Chronopharmacology of melatonin's effects in epilepsy.....	00
4.2. Melatonin and depression/anxiety comorbidities in epilepsy.....	00
5. Molecular mechanisms linking epilepsy and co-morbid anxiety and depression: the melatonin system as a potential therapeutic target.....	00
6. Conclusions and perspectives.....	00
Conflict of interest statement.....	00
References.....	00

Abbreviations:

suprachiasmatic nuclei (SCN)

hypothalamic-pituitary-adrenal (HPA)

serotonin reuptake inhibitor (SSRI)

serotonin-norepinephrine reuptake inhibitor (SNRI)

open field (OF)

elevated plus maze test (EPM)

light-dark test (LDT)

novelty suppressed feeding test (NSFT)

benzodiazepine (BZ)

hole-board test (HBT)

kainic acid (KA)

major depression disorder (MDD)

forced swim test (FST)

tail suspension test (TST)

glucocorticoid receptor (GR)

carbamazepine (CBZ)

status epilepticus (SE)

pentylentetrazole (PTZ)

N-methyl-D-aspartate (NMDA)

maximal electroshock seizure (MES)

spontaneously hypertensive rats (SHRs)

attention deficit hyperactivity disorder (ADHD)

Abstract

There is emerging evidence of the beneficial role of the melatonin system in a wide range of psychiatric and neurologic disorders, including anxiety, depression, and epilepsy. Although melatoninerbic drugs have chronobiotic and antioxidant properties that positively influence circadian rhythm resynchronization and neuroprotection in neurodegenerative disorders, studies examining the use of melatonin for epilepsy's comorbid psychiatric and neurological symptomatology are still limited. Preclinical and clinical findings on the beneficial effects of the melatonin system on anxiety, depression, and epilepsy suggest that melatoninerbic compounds might be effective in treating comorbid behavioral complications in epilepsy beyond regulation of a disturbed sleep–wake cycle.

Introduction

Secretion of the hormone melatonin is under the control of the main circadian clock, located in the suprachiasmatic nuclei (SCN) of the hypothalamus. Although the circadian synchronizing activity of melatonin is effective at specific time-points in diurnal and nocturnal species, the highest secretion of melatonin in all species, whether diurnal or nocturnal, occurs during the dark phase (peak at 2.00 a.m.), with the lowest occurring during the light phase (Arendt, 1998). Furthermore, melatonin deficit after pinealectomy was found to cause a phase shift without abolishment of the circadian rhythm of the rest–activity cycle (Cheung and McCormack, 1982).

The presence of melatonin receptors in the SCN and the pineal gland determines the bidirectional relationship between these structures. Melatonin receptor expression is under the control of light, and secretion of endogenous ligand follows a circadian rhythm (Salva and Hartley, 2012). While melatonin synthesis is regulated by the SCN clock, the hormone directly affects the activity of the SCN through activation of melatonin receptors (Pèvet et al., 2002). Therefore, treatment with drugs targeting the two main types of melatonin receptor in the SCN, namely MT₁ and MT₂ receptors, might affect the circadian regulatory mechanism.

Clinical studies have revealed that anxiety and depression are the most common behavioral complications in patients with epilepsy (Kanner and Nieto, 1999; Paradiso et al., 2001). These comorbid disorders are characterized by disturbed circadian rhythms of a number of parameters as well as dysregulation of the hypothalamic-pituitary-adrenal (HPA) axis (Maes et al., 1993; Mazarati et al., 2008; Pariante and Lightman, 2008; O'Toole et al., 2014). Furthermore, close relationships between the sleep–wake cycle and the diurnal rhythmicity of seizure activity (Quigg, 2000), as well as between seizure susceptibility and HPA axis activity (O'Toole et al., 2014), have been found.

Mostly prescribed for resynchronization of disturbed biological rhythms in a variety of pathological conditions, melatonin has low toxicity, neuroprotective effects, and a powerful antioxidant action. The chronobiotic activity of the hormone has been validated for cardiovascular, immune, digestive, and temperature regulation (Pevet and Challet, 2011). The MT₁ and MT₂ receptor agonist and serotonin-2C (5-HT_{2C}) antagonist agomelatine, developed by the pharmaceutical company Servier, was introduced as a new class of antidepressant in Europe in 2009, and has been intensively studied in different neurodegenerative disease models (Dastgheib and Moezi, 2014; Descamps et al., 2014; Stahl, 2014). Stimulation of the melatonin system was found to positively influence symptoms of depression, which is commonly linked with anxiety and a disturbed sleep–wake cycle. Experimental and clinical data revealed that while melatonin alone does not have any clear-cut antidepressant or anxiolytic action, agomelatine does exert antidepressant and anxiolytic effects because of its dual pharmacodynamic activity as a melatonin receptor agonist on the one hand and a 5-HT_{2C} receptor antagonist on the other (Bourin et al., 2004; Millan et al., 2005; Papp et al., 2006). Like melatonin, the efficacy of agomelatine is associated mostly with its capacity to correct abnormal circadian fluctuations in many physiological functions (Mairesse et al., 2013; Castanho et al., 2014).

Disturbance of the circadian rhythm of melatonin synthesis, and its relationship with changes in HPA axis activity, has been studied in patients with depression (Beck-Friis et al., 1985; Claustrat et al., 1984). Several studies have reported that melatonin deficit due to pinealectomy exacerbates epileptogenesis and is associated with neuronal loss in particular brain structures, such as the hippocampus and amygdala (De Lima et al., 2005; Janjoppi et al., 2006; Yildirim et al., 2013); moreover, a deficit of melatonin can negatively affect the HPA axis (Weidenfeld et al., 1993). Indeed, exogenous melatonin has been shown to abolish the circadian pattern of spontaneous epileptic seizures and to exert beneficial effects on seizure

frequency, neuronal damage, status epilepticus (SE)-induced oxidative stress, and behavioral complications concomitant with epilepsy (Tchekalarova et al., 2013, Petkova et al., 2014; Atanasova et al., 2013). Therefore, chronobiotic melatonergic drugs may be used to reverse two contributors to the development of comorbid psychiatric complications – depression and anxiety – desynchronizing the rhythms of a number of physiological parameters and attenuating the hyper-activated HPA axis seen in epilepsy.

The important role of the MT₁ receptor for maintenance of HPA axis activity has been confirmed recently with MT₁ receptor knockout mice, which have a blunted circadian fluctuation in diurnal corticosterone levels (Comai et al., 2015). Targeting this receptor has beneficial effects that might indirectly influence pathological hyperexcitability, oxidative stress, and imbalance of excitatory/inhibitory neurotransmission in the brain, which have been associated with disturbances in behavioral responses, memory deficit, neuronal death, and plastic changes in the hippocampus.

This review summarizes current findings on the role of the melatonergic system in neurodegenerative disorders, with a focus on anxiety, depression, and epilepsy. We hypothesize that melatonergic drugs targeting MT₁ and MT₂ receptors as well as possessing an antagonistic effect on the 5-HT_{2C} receptor subtype can be used as an effective add-on therapy in epilepsy for the treatment of comorbid anxiety and depression.

1. The melatonin system and anxiety

Although patients with panic disorder have decreased plasma melatonin at night (McIntyre et al., 1987; Cameron and Nesse, 1988), anxiogenic stimuli have been reported to evoke secretion of melatonin from the pineal gland in mice (Golombek et al., 1996). Several clinical reports have confirmed the impact of perioperative treatment with melatonin as an anxiolytic

(Caumo et al., 2007; Khezri et al., 2013; Yousaf et al., 2010). Evidence suggests that agomelatine may be considered an alternative for the treatment of anxiety disorder (Stein et al., 2008), having a comparative, or higher, efficiency than the selective serotonin reuptake inhibitor (SSRI) class, serotonin-norepinephrine reuptake inhibitor (SNRI) class, and sertraline (reviewed in De Berardis et al., 2013).

Experimental studies have revealed that melatonin exerts a modest activity in different tests for anxiety in rodents (Table 1). The first report on the efficacy of melatonin to alleviate anxiety in experimental animals was that of Golus and King on the open field (OF) test (1981). Although there is a discrepancy among reports on the effective dose and time of delivery (evening vs. morning), there is consensus that melatonin's effects are strongly dependent upon the timing of its administration (Golombek et al., 1993; 1996; Loiseau et al., 2006; Papp et al., 2006, Tian et al., 2010), suggesting a primary role for melatonin receptors in mediating the chronobiotic properties of the hormone.

Administration of this hormone to rats two hours before or during the dark phase produced a dose-dependent anxiolytic effect in the elevated plus maze (EPM) test, the light-dark test (LDT), ultrasonic vocalization, and the Vogel test (Kopp et al.; 2000; Loiseau et al., 2006; Papp et al., 2006; El Mrabet et al., 2012). In contrast, acute administration of melatonin in the morning was found to be ineffective in most anxiety tests (Golombek et al., 1993; Loiseau et al., 2006; Millan et al., 2005; Papp et al., 2006). However, agomelatine and the melatonin agonist Neu-P11 exhibited potent anxiolytic effects independent of the timing of administration (Millan et al., 2005; Papp et al., 2006, Tian et al., 2010). Although these new drugs were shown to have slightly weaker effects than the classical anxiolytics diazepam and buspirone (Tian et al., 2010), they do not produce sedation (Papp et al., 2006). The melatonin antagonist S22153 was shown to block the activities of melatonin and agomelatine when the latter were administered in the evening but not in the morning (Papp et al., 2006), confirming

the assumption that the anxiolytic effects of the two drugs are mediated by melatonin receptors. However, the lack of a time-dependent effect on anxiety with agomelatine suggests a role for 5-HT_{2C} receptor antagonism, whereas the lack of a chronobiotic activity for Neu-P11 might be due to its interaction with GABAergic (Laudon et al., 2008) or 5-HT (Tian et al., 2010) pathways. Agomelatine's dual action on melatonergic and serotonergic components determines a broader anxiolytic efficacy and lack of chronobiotic dependence compared with melatonin.

Several studies have indicated that the anti-anxiety activity of melatonin depends on the type of experimental test employed: indeed, melatonin was shown to be effective in the Vogel test, but not in the EPM test (Loiseau et al., 2006), which implicates positive responses against a 'trait', but not 'state' anxiety (Papp et al., 2006). Genetic studies on MT₁ and MT₂ receptor knockout mice confirmed the principal role of both receptors in anxiety responses (Weil et al., 2006; Comai et al., 2014; 2015). Interestingly, these knockout mice exhibited anxiety-resistant behavior in the OF test, in contrast to the anxiogenic activity shown in the novelty suppressed feeding test (NSFT) (Comai and Gobbi, 2014; Comai et al., 2015); the authors suggested that NSFT was a more specific test for validating anxiety-like behavior in genetically manipulated animals. Moreover, whereas hyperactivity of MT₁^{-/-} mice produced false-positive hypoanxiety behavior with the EPM test (Comai et al., 2015), knockout of the MT₂ receptor reflected on decision-oriented behavior (Comai and Gobbi, 2014). A previous study by the same research group supports the principal participation of the MT₂ receptor in anxiety (Ochoa-Sanchez et al., 2012). Furthermore, the MT₂-selective partial agonist UCM765 was reported to be an effective anxiolytic compound even at a low dose, and did not produce side effects, such as sedation and abuse.

Studies on how melatonin deficit produced by pinealectomy affects the level of anxiety are controversial. Some authors indicate no influence of pinealectomy with the EPM test

(Appenrodt and Schwarzberg, 2000; Juszczak et al., 1996; Karakaş et al., 2011), whereas others report increased anxiety with the burying behavior test (Bustamante-García et al., 2014), the OF test, and the EPM test (El Mrabet et al., 2012). This inconsistency could be due to differences in the testing period. Indeed, it has recently been demonstrated that anxiety levels return to normal three months after removal of the pineal gland, suggesting involvement of a compensatory, extra-pineal, melatonin system (Nenchovska et al., 2014).

A bulk of data in the literature shows that benzodiazepine (BZ) receptors are also involved in the emotional effects of melatonin, and that melatonin can modulate the GABA-BZ receptor complex. Exogenous melatonin has been demonstrated to enhance GABA-induced chloride influx and GABA turnover rate (Rosenstein and Cardinali, 1986; Rosenstein et al., 1989), as well as changing GABA-BZ binding to brain membranes (Acuña-Castroviejo et al., 1986; Niles et al., 1987). The combination of either melatonin or agomelatine with diazepam is unable to enhance the myorelaxant activity of the BZ drug, confirming the pharmacodynamic nature of their relationship (Loiseau et al., 2006), but melatonin does potentiate diazepam's anxiolytic action, as indicated by several anxiety tests on mice and rats (Guardiola-Lemaître et al., 1992; Loiseau et al., 2006), and exerts a dose-dependent effect in the conflict test in rats (Naranjo-Rodriguez et al., 2000). Furthermore, the anxiolytic activity of melatonin is antagonized by pretreatment with the BZ antagonist flumazenil in the EPM procedure in rats (Golombek et al., 1993), the hole-board test (HBT), LDT, and free-exploratory tests in mice (Pierrefiche et al., 1993; Kopp et al., 1999). In contrast, the anxiolytic profile of agomelatine seems to be different from that of BZs (Papp et al., 2006, Millan et al., 2005), and the potential role of 5-HT_{2C} receptor antagonism underlying the anxiolytic activity of the former is still uncertain and needs further study.

Melatonin restored anxiety to a normal level after application of different aversive stimuli, such as administration of lipopolysaccharide (Nava and Carta, 2001), diazinon (Ahmed et al.,

2013), or doxorubicin (Aziriova et al., 2014), and upon long-term restrain stress in mice (Kumar et al., 2014). Continuous administration of melatonin had disease-modifying effects on neuronal damage and behavioral disturbances in different models of neurodegenerative disorders: long-term melatonin treatment (4 mg/kg/day, via drinking water for 29 days) attenuated behavioral complications associated with increased motor activity and anxiety in a mouse model of ischemic stroke (Kilic et al., 2008), but repeated melatonin administration was unable to prevent kainic acid (KA)-induced epileptogenesis in Wistar rats and spontaneously hypertensive rats (SHRs), although it did exert neuroprotection in specific brain areas and restored the anxiety response during the chronic epileptic phase (Tchekalarova et al., 2013; Petkova et al., 2014).

2. Melatonin and depression

Patients with major depression disorder (MDD) are characterized by a low melatonin level at night, a disturbed HPA axis, and a phase shift in circadian melatonin and cortisol rhythms (Beck-Friis et al., 1985; Nair et al., 1984). Although agomelatine is as effective as other classic antidepressants in the acute phase of MDD, insufficient data on its efficacy for relapse prevention and its hepatotoxic side-effects hampers its use as a first-choice therapeutic for MDD (Gahr, 2014). Overall, assessment of melatonin could be effective in acute tests for the screening of antidepressants as well as in chronic models of depression. However, clinical data have not revealed clear-cut evidence that melatonin, per se, is able to prevent depressive symptoms in MDD. Indeed, long-term treatment with a combination of melatonin and low doses of the 5-HT_{1A} receptor agonist buspirone exerted a beneficial activity and few side effects in patients with acute MDD (Fava et al., 2012). It seems that the beneficial effect of melatonin treatment as an adjuvant to other antidepressants is associated with its ability to synchronize disturbed circadian rhythms in patients with depression and a deficit of melatonin

secretion (Haimov et al., 1995; Garfinkel et al., 1995; Jan et al., 2001; Mittal et al., 2010; Wade et al., 2010).

Pre-clinical and clinical data support the notion that the selective MT₁/MT₂ agonist ramelteon, which has a substantially longer half-life than melatonin, might be considered as a drug of choice for the treatment of circadian rhythm sleep disorders (Salva and Hartley, 2012). Despite its chronobiotic activity as a pure MT₁ and MT₂ receptor agonist, ramelteon is unlikely to affect depressive symptoms in MDD (Salva and Hartley, 2012). Currently approved in Europe for the treatment of MDD, agomelatine improves the sleep and clinical symptoms of depressive illness and does not have the side effects on sleep seen with other compounds in use (Arendt and Rajaratnam, 2008; Eser et al., 2007). Preclinical data has revealed that exogenous melatonin is active in routinely used screening tests for antidepressant actions, such as the forced swim test (FST) (Bourin et al., 2004; Brotto et al., 2000; Ergün et al., 2008; Micale et al., 2006; Overstreet et al., 1998; Raghavendra et al., 2000; Shaji et al., 1998; Ramírez-Rodríguez et al., 2014; Zahra et al., 2012;) and the tail suspension test (TST) (Prakhie et al., 1998; Mantovani et al., 2003; Binfarè et al., 2010; Oxenkrug et al., 2010) (Table 2). Agomelatine was reported to be more efficacious than melatonin in the FST on naïve rats and mice (Bourin et al., 2004) as well as on transgenic mice with low glucocorticoid receptor (GR) function (Barden et al., 2005). Moreover, the efficacy of melatonergic drugs has been shown to be strongly dependent upon sex hormones, species (rats vs. mice), sex (male vs. female), and treatment protocol (chronic vs. acute) (Bourin et al., 2004; Brotto et al., 2000; Paizanis et al., 2010; Raghavendra et al., 2000; Zahra et al., 2012). Indeed, chronic, but not acute, treatment regimens with pharmacological doses of melatonin and agomelatine had an anti-depressant effect in mice (Raghavendra et al., 2000; Bourin et al., 2004), whereas acute and chronic administration of agomelatine were effective in an FST on rats (Bourin et al., 2004). Unlike agomelatine, the activity of melatonin

is strongly dependent on the timing of administration, with an evening antidepressant activity for melatonin receptors and a morning activity for 5-HT_{2C} receptors (Papp et al., 2003). Clinical data has also revealed that the period of delivery of exogenous melatonin is critical for the efficacy of this hormone on the circadian rhythms of sleep and core body temperature (Arendt, 2005).

The antidepressant activity of melatonin and melatonin receptor agonists has been confirmed in a genetic animal model of depression (Flinders Sensitive Line rats) (Overstreet et al., 1998), in MT₁^{-/-} mice on an FST (Weil et al., 2006), and in the chronic mild stress model of depression in mice and rats (Kopp et al., 1999; Papp et al., 2003; Haridas et al., 2013). Furthermore, a recent study on MT₁^{-/-} mice has suggested that the MT₁ receptor is implicated in melancholic depression, and that MT₁ receptor agonists might be considered as potential antidepressant drugs with putative chronobiotic effects (Comai et al., 2015). In contrast, the role of MT₂ receptors in depression is not clear, although there are some indications that the antidepressant-like activity of luzindole is mediated through this receptor sub-type (Sumay et al., 2005).

Melatonin was less effective than fluoxetine in preventing chronic mild-stress-induced complications in the behavior of C3H/He mice (Kopp et al., 1999). Agomelatine had an antidepressant effect in a rat model of prenatal restraint stress, suggesting that it might be considered a “disease-dependent drug” (Morley-Fletcher et al., 2011). The combination of agomelatine with 5-HT agonists/antagonists was more efficacious than the effects of melatonin alone. While the putative role of 5-HT_{2C} receptor blockade in the mechanisms underlying the antidepressant potency of agomelatine is supported by experimental and clinical data, the mechanism of melatonin’s antidepressant activity remains obscure. The antidepressant effect of melatonin in BALB/c mice was suppressed by pretreatment with a 5-HT_{2A}/5-HT_{2C} agonist and BZ receptor antagonist, suggesting involvement of 5-HT and BZ

receptors (Raghavendra et al., 2000; Micale et al., 2006). On the other hand, experimental data has indicated that the effect of melatonin in the TST might be mediated by an interaction with dopamine D₁ and D₂ receptors, NMDA receptors, the L-arginine-NO pathway, and the BZ pathway (Binfarè et al., 2010; Mantovani et al., 2003; Guardiola-Lemaître et al., 1992). The above-mentioned results reveal the complex mechanisms underlying melatonin's antidepressant activity, which need to be explored in more detail.

3. Melatonin and epilepsy

Melatonin's antioxidant activity and low toxicity makes it attractive to clinicians as a potential adjuvant for epilepsy therapy. Furthermore, studies have reported that the primary beneficial effect of melatonin on seizure activity is due to its ability to synchronize disturbed circadian rhythms in epileptic patients (Fauteck et al., 1999; Peled et al., 2001). Comorbidity of a sleep disorder with epilepsy creates difficulties in choosing an anti-epileptic drug (AED). Indeed, while lamotrigine, levetiracetam, valproate, and gabapentine have been reported to lack or exert few negative effects, other agents such as BZs, barbiturates, and phenytoin aggravate sleep architecture in individuals with epilepsy (reviewed in: Bazil et al., 2003). In line with this, melatonin and melatonin receptor agonists may be suitable for use as add-on treatments with routine AEDs. Indeed, melatonin alleviates sleep disturbances of different etiologies in epileptic children and adults (Coppola et al., 2004; Fauteck et al., 1999; Peled et al., 2001). The improvement in sleep parameters is accompanied by an attenuation of seizure activity in young patients with intractable epilepsy (Elkhayat et al., 2010) and in patients on valproate (Gupta et al., 2005), but melatonin produces an ambiguous, dual effect in epileptic patients with mental complications (Coppola et al., 2004). Furthermore, clinical reports have shown the beneficial effect of melatonin treatment on seizure activity during the day (Goldberg-Stern et al., 2012) and the night (Peled et al., 2001) in patients with intractable epilepsy, a pathology

characterized by an increased post-seizure salivary melatonin level (Bazil et al., 2003). High melatonin levels were also detected in unmedicated patients with complex partial epilepsy, and might represent a compensatory feedback mechanism against the severe epileptic state (Graham et al. 1995). However, increased night-time plasma melatonin has been associated with a higher incidence of seizure attacks at night (Sandyk et al., 1992).

Another important advantage of melatonin as an add-on option for epilepsy is associated with its potent antioxidant properties. Clinical data has shown that melatonin has an antioxidant activity in epileptic patients on carbamazepine (CBZ) (Gupta et al., 2004, 2006). Increased oxidative stress and disturbance of the endogenous antioxidant system can aggravate seizure activity (Yüksel et al., 2000). On the other hand, data has shown that the metabolites of several AEDs behave as pro-oxidant agents with toxic effects (Graf et al., 1998; Niketic et al., 1995; Yüksel et al., 2000). Experiments have confirmed the potent antioxidant activity of melatonin in different pathological states: chronic melatonin infusion (10 mg/kg/day), as well as a cumulative dose of 10 mg/kg, suppressed KA-induced oxidative stress in the hippocampus (Chung and Han, 2003) without preventing the development of SE (Atanasova et al., 2013).

Melatonin has been reported to have an anticonvulsant action in many models of acute seizures, such as those produced by the administration of pentylenetetrazole (PTZ), picrotoxin, bicuculline, pilocarpine, L-cysteine, kainate, 3-mercaptopropionic acid, quinolinate, glutamate (Glu), strychnine, N-methyl-D-aspartate (NMDA), or penicillin, as well as in the maximal electroshock seizure (MES) test in rats, mice, gerbils, and hamsters (Table 3). The anticonvulsant and neuroprotective effects of melatonin have been verified in the chronic amygdala kindling model of partial epilepsy in rats (Mevissen et al., 1998) and the KA model of temporal lobe epilepsy (Tchekalarova et al. 2013; Petkova et al., 2014). Pinealectomy facilitates epileptogenesis in some species and increases seizure activity (De

Lima et al., 2005; Janjoppi et al., 2006; Silva de Lacerda et al., 2007; Yildirim et al., 2013), and melatonin has been reported to either reverse (Yildirim et al., 2013) or to be ineffective (De Lima et al., 2005) in this pinealectomy-induced enhancement of seizure susceptibility.

In addition to the MT receptor-mediated effects of melatonin in seizure models, a number of studies have demonstrated the participation also of GABAergic, 5-HT-ergic, and NO/L-arginine pathways (Bikjdaouene et al., 2003; Dastgheib and Moezi, 2014; Golombek et al., 1992; Ray et al., 2004; Yahyavi-Firouz-Abadi et al., 2007). Melatonin potentiates the effect of several AEDs, such as carbamazepine and phenobarbital (Borowicz et al., 1999; Forcelli et al., 2013), but fails to protect neonatal rats against PTZ-induced seizures, even at the high dose of 80 mg/kg (Forcelli et al., 2013). Although chronic melatonin infusion had an antioxidant effect in Wistar rats and SHRs with KA-induced SE, it was ineffective against the development of SE (Atanasova et al., 2013); however, it did partially blunt pilocarpine-induced SE in pinealectomized rats (De Lima et al., 2005). Moreover, a few reports have raised the point that melatonin can act as an endogenous proconvulsant (Stewart and Leung, 2003; Yehudi and Mostofsky, 1993). These contrasting effects might be associated with Mel_{1b} receptor-mediated suppression of GABA_A receptors in CA1 pyramidal cells (Stewart and Leung, 2003) and involvement of the dopaminergic pathway (Yehudi and Mostofsky, 1993).

The first study considering the efficacy of agomelatine in different seizure models in female Swiss mice was published by Aguiar et al. (2012). Similarly with previously reported results on the anticonvulsant effects of melatonin at doses ranging from 40 to 80 mg/kg in PTZ-induced seizures in a variety of species (Bikjdaouene et al., 2003; Moezi et al., 2011; Solmaz et al., 2009; Yahyavi-Firouz-Abadi et al., 2007), the authors showed that agomelatine was active at doses of 25 and 50 mg/kg. Agomelatine was also effective against pilocarpine-induced SE at a dose of 75 mg/kg (Aguiar et al., 2012), while melatonin had an anticonvulsant effect only after repeated injections in this model of epilepsy (Costa-Lotufu et

al., 2002; De Lima et al., 2011). The lack of activity of agomelatine against other convulsants, such as picrotoxin and strychnine, as well as in the MES test, suggests the presence of more-specific mechanisms of action, which need further exploration. The dose-dependent anticonvulsant effect of acute agomelatine administration against PTZ seizure threshold in mice was recently confirmed by Dastgheib and Moezi (2014). However, agomelatine was ineffective in a chronic treatment protocol, a result possibly due to agomelatine-induced receptor desensitization or internalization. In line with this, the receptor-mediated melatonin effect was also found to involve a fine-tuning mechanism of desensitization or internalization, which was strictly dependent upon the treatment protocol (Gerdin et al., 2004; Delagrangue and Guardiola-Lemaitre, 1997; Witt-Enderby et al., 2003). Moreover, the anticonvulsant efficacy of agomelatine against i.v.-injected PTZ in mice was reported to involve iNOS or nNOS induction, because selective iNOS and nNOS inhibitors attenuated the effect of agomelatine (Dastgheib and Moezi, 2014). These results concurred with previous findings that the NO/L-arginine pathway was involved in the anticonvulsant effect of melatonin against clonic seizures induced by i.v.-infused PTZ (Bikjdaouene et al., 2003; Yahyavi-Firouz-Abadi et al., 2006).

In conclusion, clinical and experimental findings showing the beneficial effect of melatonin on disturbed circadian rhythms and oxidative stress in epileptic conditions suggest that the targeting of the pathways induced by this hormone may be used to treat epilepsy-related complications. However, because 5-HT_{2C} receptors are also involved in the regulation of seizure susceptibility (Applegate and Tecott, 1998; Isaak, 2005; Upton et al., 1998), further studies on models of chronic epilepsy are required to gain a better understanding of the mechanisms underlying agomelatine's anti-seizure effect.

4.1. Chronopharmacology of melatonin's effects on epilepsy

Few studies have focused on the chronopharmacology of melatonin in seizure tests and models of epilepsy. Moreover, data is controversial: indeed, exogenous melatonin has been reported to produce an anticonvulsant effect that was either strongly time-dependent (Costa-Lotufo et al., 2002; Golombek et al., 1992; Lapin et al., 1998) or independent from the time of day of administration (Champney et al., 1996; Mevissen et al., 1998). Exogenous melatonin and the secretion of the endogenous hormone at night can both protect neuronal cultures from apoptosis promoted by KA-induced oxidative stress (Manev et al., 1996). Melatonin produced a diurnal variation in susceptibility to PTZ- and 3-MP-induced seizures in hamsters, with the lowest threshold needed to evoke clonic seizures found at night (Golombek et al., 1992). Melatonin administered at a dose of 50 mg/kg i.p. had the greatest effect in the early evening, when the level of endogenous hormone is expected to be the most elevated (Golombek et al., 1992). Melatonin activity seems to be influenced by central BZ antagonism, because Ro 15-1788 and flumazenil blocks its effects. The circadian rhythm of seizure threshold has been documented for different seizure models, including PTZ (Yehuda and Mostofsky, 1993), pilocarpine (Stewart, 2001), and MES test (Löscher et al., 1996). Melatonin was ineffective during the day in rats, but decreased PTZ seizure threshold at night (Yehuda and Mostofsky, 1993).

Clinical and experimental data has revealed the circadian rhythm of seizure activity in patients and in different models of epilepsy (Arida et al., 1999; Raedt et al., 2009; Shouse et al., 1996; Stewart and Leung, 2003). Indeed, there is a prevalence of epileptiform activity during sleep (Asano et al., 2007; Ferrillo et al., 2000; Staba et al., 2002). Rodents are nocturnal animals, and a higher frequency of EEG and video-recorded motor seizures have been monitored in different models of epilepsy when animals are inactive during the light phase (Arida et al., 1999; Bertram et al., 1994; Quigg, 2000; Raedt et al., 2009; Tchekalarova et al., 2013).

Similar higher seizure susceptibility during the inactive period was found in rats subjected to amygdala kindling-induced epileptogenesis (Weiss et al., 1993). Recently, we demonstrated that the administration of melatonin flattens the circadian periodicity of spontaneous seizures in Wistar rats and SHRs with KA-induced temporal lobe epilepsy (Tchekalarova et al., 2013, Petkova et al., 2014). The effect was greater in SHRs, and possibly related to decreased seizure frequency during the light phase.

The circadian periodicity of endogenous melatonin secretion might also influence the diurnal pattern of seizures. Yalín et al. (2006) reported that melatonin level and its circadian rhythm do not differ between patients with nocturnal or diurnal complex partial seizures. This finding suggests that seizure periodicity does not depend on fluctuations in melatonin release. Melatonin receptor expression follows a circadian pattern and depends on exposure to either light or melatonin. Furthermore, the administration of melatonin produces a phase shift in circadian fluctuations. It is accepted that this chronobiotic activity of the hormone is due to the specific and divergent roles of MT₁ and MT₂ receptors in the central biological clock, located in the SCN at the anterior hypothalamus (Liu et al., 1997).

Studies have demonstrated that humans and experimental animals with epilepsy are characterized by shifted or disrupted circadian rhythms in cortisol release (Bazil et al., 2003), temperature (Quigg, 2000), motor activity (Stewart and Leung, 2003), and sleep architecture, for example decreased REM sleep (Bastlund et al., 2005; Manni and Terzaghi, 2010). Fluctuations of the melatonin level in the sleep–wake cycle have been shown to correlate with a propensity for epileptic activity in patients with complex partial epilepsy (Graham et al. 1995) or following seizures (Bazil et al., 2003), suggesting a compensatory, melatonin-mediated feedback mechanism against severe epilepsy. Studies on the circadian rhythm of melatonin release during epilepsy are missing. Work is needed to understand whether and

how melatonin release and the expression of MT receptors in various brain structures are altered during or related to the circadian pattern of spontaneous seizures.

4.2. Melatonin and depression/anxiety comorbidities in epilepsy

Behavioral complications, such as depression, anxiety, memory deficit, and attention deficit hyperactivity disorder (ADHD), are common concomitances in epilepsy. The predisposition of patients with psychiatric complications for seizures and of patients with epilepsy to develop behavioral complications are both well documented (Adelöw et al., 2012; Forsgren and Nystrom, 1999; Hesdorffer et al., 2012). Although data favors the hypothesis that epilepsy and its behavioral comorbidities have a common underlying mechanism, and that morphological, biochemical, and behavioral changes caused by one disorder may provoke another, the exact pathogenesis of behavioral comorbidities in epilepsy remains obscure, and there is a lack of standard screening and treatment approaches (Ott et al., 2003). Over the last decade, several laboratories have validated animal models of comorbidity, such as depression and ADHD (Koh et al., 2007; Ma and Leung, 2004; Mazarati et al., 2008; Pineda et al., 2010; 2014). We have recently reported on strain-related behavioral differences, using the KA-induced epilepsy model, which is also considered a model of depression comorbidity (Tchekalarova et al., 2010). The intact SHR is characterized by typical behavioral, electrophysiological, and biochemical abnormalities – which are also evident in epileptic rats – such as attention deficit, impulsiveness, deficient sustained attention, and monoamine deficits in the frontal cortex and hippocampus, suggesting that this strain might be used also as a relevant model of ADHD comorbidity (Tchekalarova, 2014). In addition, we recently demonstrated that long-term treatment with melatonin for KA-induced epileptogenesis produces different effects in Wistar rats and SHRs; the finding supported the hypothesis that

the underlying mechanisms of behavioral comorbidities in epilepsy differ in these two rat strains (Tchekalarova et al., 2013; Petkova et al., 2014). Indeed, while seizure frequency in epileptic SHRs remained significantly decreased even after discontinuation of melatonin, in Wistar rats the anticonvulsant effect of melatonin was evident only during the treatment period. In addition, whereas the administration of melatonin to Wistar rats produced a beneficial effect on behavioral and biochemical complications, such as hyperactivity, anxiety, depression, memory deficit, and low hippocampal 5-HT (Tchekalarova et al., 2014), long-term treatment failed to positively affect the behavioral and memory deficits or improve the disturbed diurnal behavioral rhythms in epileptic SHRs (Petkova et al., 2014). However, melatonin was neuroprotective in the CA1 hippocampal area and the piriform cortex in both rat strains, suggesting a positive relationship between anti-seizure efficacy and neuroprotection.

So far, the efficacy of agomelatine for depression, anxiety, memory deficit, and sleep disturbances has been proven in humans and in animal models (reviewed in Vimala et al., 2014). Experimental and clinical studies on the anticonvulsant potency of acute agomelatine administration on several seizure tests (Aguilar et al., 2012; Dastgheib and Moezi, 2014) and the positive effect on behavioral disturbances suggest that this agent might have beneficial effects also in epilepsy for comorbid depression, anxiety, and sleep disturbances.

4. Molecular mechanisms linking epilepsy and comorbid anxiety and depression: the potential of targeting the melatonin system

Basic molecular mechanisms of the melatonin system that might be considered as possible targets:

- i) Melatonin-mediated restoration of a disrupted hypothalamic-pituitary-adrenal axis – Clinical data and experimental models have shown that the HPA axis is hyperexcited in

epilepsy (Culebras et al., 1987; Mazarati et al., 2009; O'Toole et al., 2014) and in depression (Cai et al., 2013; Chen et al., 2015) and that melatonin can correct glucocorticoid-induced HPA axis disruption (Konakchieva et al., 1998). Because stress responses are regulated by the hypothalamic paraventricular nucleus (Laryea et al., 2013), and MT₂ receptors are present in this area (Lacoste et al., 2015), we may speculate that melatonin could correct HPA axis disruption by acting on these receptors.

ii) Melatonin and GABA neurotransmission – Enhancing GABAergic responses in the brain is a common therapeutic approach for epilepsy, anxiety, and other neurological conditions. Melatonin exerts its activity either through a non-receptor-mediated mechanism (Li et al. 2001) or by binding to the GABA_A receptor (Niles and Peace, 1990), which enhances its affinity for the agonist (Wu et al. 1999). Furthermore, enhancement or attenuation of GABA_A receptor-mediated currents strongly depends on (MT₁ or MT₂) receptor activation (Wan et al. 1999). Because of the marked side effects of GABA-mimetic drugs, there is a tendency in clinical practice to replace them with melatonin agents.

iii) Melatonin and the serotonin receptor – MT₂ receptors are involved in the pathophysiology and pharmacology of sleep disorders, anxiety, and depression, and selective MT₂ receptor agonists have hypnotic and anxiolytic properties (Comai et al., 2014). Serotonin disturbances are involved in depressive (Meltzer, 1989) and epileptic (Richerson et al., 2011) conditions. A novel finding is the presence of MT₂/5-HT heteromers in the brain (Kamal et al., 2015). Melatonin could act through this heteromer to modulate epileptic seizures and depressive-like comorbidity. Therefore, this heteromer could further explain similar actions between melatonin and agomelatine.

5. Conclusions and future perspectives

Disturbance of the circadian rhythms of a number of physiological functions is relatively frequent in many neurodegenerative disorders, including depression, anxiety disorder, and epilepsy (Gorwood, 2012; Quigg, 2000; Turek, 2007). A bidirectional link between circadian rhythm and these disorders has been suggested (Salva and Hartley, 2012; Yalyn et al., 2006; Yousaf et al., 2010). Agents that synchronize circadian fluctuations are considered to possess chronobiotic activity (Pévet et al., 2002).

The rationale of using drugs that stimulate prolonged melatonin release (circadian), such as ramelteon (an MT₁ and MT₂ receptor agonist) and agomelatine (a melatonin receptor agonist and a 5HT_{2C} receptor antagonist), as adjuvants to anticonvulsant therapy in epilepsy is two-fold. First, treatment with melatonin-related drugs in epilepsy can have beneficial effects on sleep cycle synchronization, thereby exerting indirect improvements in comorbid depression and anxiety, as well as controlling seizure frequency. The presence of melatonin receptors in the SCN suggests that circulating melatonin agonists affect circadian rhythm regulation. Clinical reports have revealed that although melatonin can synchronize circadian rhythm, limiting therapeutic targeting only to MT₁ and MT₂ receptors is not sufficient to treat depressive symptoms (Dalton et al., 2000; Serfaty et al., 2010). A combination of chronobiotic (melatonin receptors) and antidepressant (5HT_{2C} antagonism) effects is proven to be a more effective therapeutic strategy for depression (Kasper and Hajak, 2013). Studies indicated that agomelatine has a beneficial influence on a group of patients with sleep problems (Kasper and Hajak, 2013; Sansone and Sansone, 2011). Besides, studies have documented general improvements in the quality of life of epilepsy patients on melatonin add-on therapy, which could be related to melatonin's chronobiotic activity (Jones et al., 2005). Second, melatonin is proven to possess neuroprotective (Manev et al., 1996; Mevissen et al., 1998; Tchekalarova et al., 2013) and antioxidant activities through scavenging of free radicals and potentiation of the activity of endogenous antioxidant enzymes (Yonei et al., 2009). Most AEDs directly target the receptors of glutamate and/or GABA, thereby causing a number of side effects. Melatonin appears to diminish the glutamate-targeting effect, while enhancing GABA activity through indirect fine-tuning mechanisms targeting melatonin receptors, promoting an inhibition of seizures with fewer side effects. Over recent years, agomelatine has become the focus of interest because of its combined chronobiotic (melatonin receptor agonism) and antidepressant (5-HT_{2C} receptor antagonism) effects. The few experimental

studies reporting an anticonvulsant effect for agomelatine in acute seizure tests give encouraging signs and suggest that this drug represents a promising therapeutic tool for the targeting of circadian desynchronization, oxidative stress, and comorbid depression and anxiety in epilepsy.

Conflict of interest statement

The authors have no conflicts of interest to declare.

Acknowledgements

We sincerely thank Michael Latronico for his expert assistance in the editing of the manuscript. The work was funded by...

References

- Acuña-Castroviejo, D., Rosenstein, R.E., Romeo, H.E., Cardinali, D.P., 1986. Changes in gamma-aminobutyric acid high affinity binding to cerebral cortex membranes after pinealectomy or melatonin administration to rats. *Neuroendocrinology* 43, 24–31.
- Adelöw, C., Andersson, T., Ahlbom, A., Tomson, T., 2012. Hospitalization for psychiatric disorders before and after onset of unprovoked seizures/epilepsy. *Neurology* 78, 396–401.
- Aguiar C.C., Almeida, A.B., Araújo, P.V., Vasconcelos, G.S., Chaves, E.M., do Vale, O.C., Macêdo, D.S., de Sousa, F.C., Viana, G.S., Vasconcelos, S.M., 2012. Anticonvulsant effects of agomelatine in mice. *Epilepsy & Behavior* 24, 324-328.
- Ahmed, M.A., Ahmed, H.I., El-Morsy, E.M., 2013. Melatonin protects against diazinon-induced neurobehavioral changes in rats. *Neurochemistry Research* 38, 2227-2236.
- Applegate, C.D., Tecott, L.H., 1998. Global increases in seizure susceptibility in mice lacking 5-HT_{2C} receptors: a behavioral analysis. *Experimental Neurology* 154, 522-530.
- Appenrodt, E., Schwarzberg, H., 2000. Central vasopressin administration failed to influence anxiety behavior after pinealectomy in rats. *Physiology and Behavior* 68, 735-739.
- Arendt, J. 2005. Melatonin: characteristics, concerns, and prospects. *Journal of Biological Rhythms* 20, 291–303.
- Arendt, J., Rajaratnam, S.M., 2008. Melatonin and its agonists: an update. *British Journal of Psychiatry* 193, 267-269.
- Arida, R.M., Scorza, F.A., Peres, C.A., Cavalheiro, E.A., 1999. The course of untreated seizures in the pilocarpine model of epilepsy. *Epilepsy Research*, 34, 99-107.

Asano, E., Mihaylova, T., Juhász, C., Sood, S., Chugani, H.T., 2007. Effect of sleep on interictal spikes and distribution of sleep spindles on electrocorticography in children with focal epilepsy. *Clinical Neurophysiology* 118, 1360-1368.

Aziriova, S., Repova Bednarova, K., Krajcirovicova, K., Hrenak, J., Rajkovicova, R., Arendasova, K., Kamodyova, N., Celec, P., Zorad, S., Adamcova, M., Paulis, L., Simko, F., 2014. Doxorubicin-induced behavioral disturbances in rats: protective effect of melatonin and captopril. *Pharmacology Biochemistry and Behavior* 124, 284-289.

Atanasova, M., Petkova, Zl., Pechlivanova, D., Dragomirova, P., Blazhev, Al., Tchekalarova, J., 2013. Strain differences in the effect of long-term treatment with melatonin on kainic acid-induced status epilepticus, oxidative stress and the expression of heat shock proteins. *Pharmacology Biochemistry and Behaviour* 111, 44-50.

Barden, N., Shink, E., Labbé, M., Vacher, R., Rochford, J., Mocaër, E., 2005. Antidepressant action of agomelatine (S 20098) in a transgenic mouse model. *Progress in Neuropsychopharmacology and Biological Psychiatry* 29, 908-916.

Bastlund, J.F., Jennum, P., Mohapel, P., Penschuck, S., Watson, W.P., 2005. Spontaneous epileptic rats show changes in sleep architecture and hypothalamic pathology. *Epilepsia* 46, 934-938.

Bazil, C.W., 2003. Effects of antiepileptic drugs on sleep structure : are all drugs equal? *CNS Drugs* 17, 719-728.

Beck-Friis, J., Ljunggren, J.G., Thorén, M., von Rosen, D., Kjellman, B.F., Wetterberg, L., 1985. Melatonin, cortisol and ACTH in patients with major depressive disorder and healthy humans with special reference to the outcome of the dexamethasone suppression test. *Psychoneuroendocrinology* 10, 173-186.

Bertram, E.H., Cornett, J.F., 1994. The evolution of a rat model of chronic spontaneous limbic seizures. *Brain Research* 661, 157-162.

Bikjdaouene, L., Escames, G., León, J., Ferrer, J.M., Khaldy, H., Vives, F., Acuña-Castroviejo, D., 2003. Changes in brain amino acids and nitric oxide after melatonin administration in rats with pentylenetetrazole-induced seizures. *Journal of Pineal Research* 35, 54-60.

Binfaré, R.W., Mantovani, M., Budni, J., Santos, A.R.S., Rodrigues, A.L.S., 2010. Involvement of dopamine receptors in the antidepressant-like effect of melatonin in the tail suspension test. *European Journal of Pharmacology* 638, 78–83.

Borowicz, K.K., Kamiński, R., Gasior, M., Kleinrok, Z., Czuczwar, S.J., 1999. Influence of melatonin upon the protective action of conventional anti-epileptic drugs against maximal electroshock in mice. *European Neuropsychopharmacology* 9, 185-190.

Boulle, F., Massart, R., Stragier, E., Paizanis, E., Zaidan, L., Marday, S., Gabriel, C., Mocaer, E., Mongeau, R., Lanfumey, L., 2014. Hippocampal and behavioral dysfunctions in a mouse model of environmental stress: normalization by agomelatine. *Translational Psychiatry* 4, e485.

Bourin, M., Mocaër, E., Porsolt, R., 2004. Antidepressant-like activity of S 20098 (agomelatine) in the forced swimming test in rodents: involvement of melatonin and serotonin receptors. *Journal of Psychiatry Neuroscience* 29, 126-133.

Brotto, L.A., Barr, A.M., Gorzalka, B.B., 2000. Sex differences in forced-swim and open-field test behaviours after chronic administration of melatonin. *European Journal of Pharmacology* 402, 87-93.

- Bustamante-García, R., Lira-Rocha, A.S., Espejo-González, O., Gómez-Martínez, A.E., Picazo, O., 2014. Anxiolytic-like effects of a new 1-N substituted analog of melatonin in pinealectomized rats. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 51, 133–139.
- Cai, L., Li, R., Wu, Q.Q., Wu, T.N., 2013. Effect of hesperidin on behavior and HPA axis of rat model of chronic stress-induced depression. *Zhongguo Zhong Yao Za Zhi* 38, 229-233.
- Kamal, M., Gbahou, F., Guillaume, J.L., Daulat, A.M., Benleulmi-Chaachoua, A., Luka, M., Chen, P., Kalbasi, A., Baroncini, M., Mannoury, C., Millan, M.J., Prevot, V., Delagrang, P., Jockers, R., 2015. Convergence of Melatonin and Serotonin (5-HT) Signaling at MT2/5-HT2C Receptor Heteromers. *Journal of Biological Chemistry* 290, 11537-11546.
- Cameron, O.G., Nesse, R.M., 1988. Systemic hormonal and physiological abnormalities in anxiety disorders. *Psychoneuroendocrinology* 13, 287-307.
- Castanho, A., Bothorel, B., Seguin, L., Mocaër, E., Pévet, P., 2014. Like melatonin, agomelatine (S20098) increases the amplitude of oscillations of two clock outputs: melatonin and temperature rhythms. *Chronobiology International* 31, 371-381.
- Caumo, W., Torres, F., Moreira, N.L. Jr, Auzani, J.A., Monteiro, C.A., Londero, G., Ribeiro, D.F., Hidalgo, M.P., 2007. The clinical impact of preoperative melatonin on postoperative outcomes in patients undergoing abdominal hysterectomy. *Anesthetic Analgetics* 105, 1263-1271.
- Champney, T.H., Hanneman, W.H., Legare, M.E., Appel, K., 1996. Acute and chronic effects of melatonin as an anticonvulsant in male gerbils. *Journal of Pineal Research* 20, 79-83.
- Cheung, P.W, McCormack, C.E., 1982. Failure of pinealectomy or melatonin to alter circadian activity rhythm of the rat. *Americal Journal of Physiology* 242, R261-R264.

Chung, S.Y., Han, S.H., 2003. Melatonin attenuates kainic acid-induced hippocampal neurodegeneration and oxidative stress through microglial inhibition. *Journal of Pineal Research* 34, 95-102.

Claustrat, B., Chazot, G., Brun, J., Jordan, D., Sassolas, G., 1984. A chronobiological study of melatonin and cortisol secretion in depressed subjects: plasma melatonin, a biochemical marker in major depression. *Biological Psychiatry* 19, 1215-1228.

Comai, S., Gobbi, G., 2014. Unveiling the role of melatonin MT2 receptors in sleep, anxiety and other neuropsychiatric diseases: a novel target in psychopharmacology. *Journal of Psychiatry and Neuroscience* 39, 6-21.

Comai, S., Ochoa-Sanchez, R., Dominguez-Lopez, S., Bambico, F.R., Gobbi, G., 2015. Melancholic-Like Behaviors and Circadian Neurobiological Abnormalities in Melatonin MT1 Receptor Knockout Mice. *International Journal of Neuropsychopharmacology* 18, 1-10.

Coppola, G., Iervolino, G., Mastrosimone, M., Torre, G., Ruiu, F., Pascotto, A., 2004. Melatonin in wake-sleep disorders in children, adolescents and young adults with mental retardation with or without epilepsy: a double-blind, cross-over, placebo-controlled trial. *Brain and Development* 26, 373-376.

Costa-Lotufo, L.V., Fonteles, M.M., Lima, I.S., de Oliveira, A.A., Nascimento, V.S., de Bruin, V.M., Viana, G.S., 2002. Attenuating effects of melatonin on pilocarpine-induced seizures in rats. *Comparative Biochemistry and Physiology C Toxicology & Pharmacology* 131, 521-529.

Culebras, A., Miller, M., Bertram, L., Koch, J., 1987. Differential response of growth hormone, cortisol, and prolactin to seizures and to stress. *Epilepsia* 28, 564-570.

Dalton, E.J., Rotondi, D., Levitan, R.D., Kennedy, S.H., Brown, G.M., 2000. Use of slow-release melatonin in treatment-resistant depression. *Journal of Psychiatry and Neuroscience* 25, 48-52.

Dastgheib, M., Moezi, L., 2014. Acute and chronic effects of agomelatine on intravenous penthylene-tetrazol-induced seizure in mice and the probable role of nitric oxide. *European Journal of Pharmacology* 736, 10-15.

De Berardis, D., Marini, S., Fornaro, M., Srinivasan, V., Iasevoli, F., Tomasetti, C., Valchera, A., Perna, G., Quera-Salva, M.A., Martinotti, G., di Giannantonio, M., 2013. The melatonergic system in mood and anxiety disorders and the role of agomelatine: implications for clinical practice. *International Journal of Molecular Science* 14, 12458-12383.

Delagrang, P., Guardiola-Lemaitre, B., 1997. Melatonin, its receptors, and relationships with biological rhythm disorders. *Clinical Neuropharmacology* 20, 482-510.

De Lima, E., Soares, J.M. Jr., del Carmen, Sanabria Garrido, Y., Gomes, V. S., Priel, M.R., Chada Baracat, E., Cavaleiro A.E., da Graça Naffah-Mazzacoratti, M., Amado, D., 2005. Effects of pinealectomy and the treatment with melatonin on the temporal lobe epilepsy in rats. *Brain Research* 1043, 24-31.

De Lima, E., Cabral, F.R., Cavaleiro, E.A., Naffah-Mazzacoratti, M.G., Amado, D., 2011. Melatonin administration after pilocarpine-induced status epilepticus: a new way to prevent or attenuate postlesion epilepsy? *Epilepsy & Behavior* 20, 607-612.

Descamps, A., Rousset, C., Dugua, H., Debilly, G., Delagrang, P., Cespuglio, R., 2014. Agomelatine restores a physiological response to stress in the aged rat. *Neuroscience Letters* 566, 257-262.

Elkhatat, H.A., Hassanein, S.M., Tomoum, H.Y., Abd-Elhamid, I.A., Asaad, T., Elwakkad, A.S., 2010. Melatonin and sleep-related problems in children with intractable epilepsy. *Pediatric Neurology* 42, 249-254.

El Mrabet, F.H., Ouakki, S., Mesfioui, A., El Hessni, A., Ouichou, A., 2012. Pinealectomy and Exogenous Melatonin Regulate Anxiety-Like and Depressive-Like Behaviors in Male and Female Wistar Rats. *Neuroscience & Medicine* 3, 1-10.

Ergün, Y., Orhan, F.O., Karaaslan, M.F., 2008. Combination therapy of imipramine and melatonin: additive antidepressant effect in mouse forced swimming test. *European Journal of Pharmacology* 591, 159-163.

Eser, D., Baghai, T.C., Möller, H.J., 2007. Evidence of agomelatine's antidepressant efficacy: the key points. *International Clinical Psychopharmacology* 22 (Suppl. 2), S15–S19.

Fauteck, J., Schmidt, H., Lerch, A., Kurlemann, G., Wittkowski, W., 1999. Melatonin in epilepsy: first results of replacement therapy and first clinical results. *Biological Signals Receptors* 8, 105-110.

Fava, M., Targum, S.D., Nierenberg, A., Bleicher, L.S., Carter, T., Wedel, P., Henc, R., Gagee, F., Barlow, C., 2012. An exploratory study of combination buspirone and melatonin SR in Major Depressive Disorder (MDD): A possible role for neurogenesis in drug discovery. *Journal of Psychiatric Research* 46, 1553–1563.

Ferrillo, F., Beelke, M., Nobili, L., 2000. Sleep EEG synchronization mechanisms and activation of interictal epileptic spikes. *Clinical Neurophysiology* 111, Suppl 2, S65-73.

Forcelli, P.A., Soper, C., Duckles, A., Gale, K., Kondratyev, A., 2013. Melatonin potentiates the anticonvulsant action of phenobarbital in neonatal rats. *Epilepsy Research* 107, 217-223.

Forsgren, L., Nystrom, L., 1999. An incident case referent study of epileptic seizures in adults. *Epilepsy Research* 6, 66-81.

Haimov, I., Lavie, P., Laudon, M., Herer, P., Vigder, C., Zisapel, N., 1995. Melatonin replacement therapy of elderly insomniacs. *Sleep*, 18, 598-603.

Haridas, S., Kumar, M., Manda, K., 2013. Melatonin ameliorates chronic mild stress induced behavioral dysfunctions in mice. *Physiology and Behavior* 119, 201-207.

Hesdorffer, D.C., Ishihara, L., Mynepalli, L., Webb, D.J., Weil, J., Hauser, W.A., 2012. Epilepsy, suicidality, and psychiatric disorders: a bidirectional association. *Annual Neurology* 72, 184–191.

Isaac. M., 2005. Serotonergic 5-HT_{2C} receptors as a potential therapeutic target for the design antiepileptic drugs. *Current Topics in Medical Chemistry* 5, 59-67.

Garfinkel, D., Laudon, M., Nof, D., Zisapel, N., 1995. Improvement of sleep quality in elderly people by controlled-release melatonin. *Lancet* 346, 541-544.

Gahr, M., 2014. Agomelatine in the Treatment of Major Depressive Disorder: An Assessment of Benefits and Risks. *Current Neuropharmacology* 12, 287-398.

Gerdin, M.J., Masana, M.I., Rivera-Bermúdez, M.A., Hudson, R.L., Earnest, D.J., Gillett, M.U., Dubocovich, M.L., 2004. Melatonin desensitizes endogenous MT₂ melatonin receptors in the rat suprachiasmatic nucleus: relevance for defining the periods of sensitivity of the mammalian circadian clock to melatonin. *FASEB Journal* 18, 1646-1656.

Goldberg-Stern, H., Oren, H., Peled, N., Garty, B.Z., 2012. Effect of melatonin on seizure frequency in intractable epilepsy: a pilot study. *Journal of Child Neurology* 27, 1524-1528.

Golombek, D.A., Duque D.F., De Brito Sánchez, M.G, Burin, L., Cardinali, D.P., 1992. Time-dependent anticonvulsant activity of melatonin in hamsters. *European Journal of Pharmacology* 210, 253-258.

Golombek, D.A., Martini, M., Cardinali, D.P., 1993. Melatonin as an anxiolytic in rats: time dependence and interaction with the central GABAergic system. *European Journal of Pharmacology* 237, 231-236.

Golombek, D.A., Pévet, P., Cardinali, D.P., 1996. Melatonin effects on behavior: possible mediation by the central GABAergic system. *Neuroscience and Biobehavioral Review* 20, 403-412.

Golus, P., King, M.G., 1981. The effects of melatonin on open field behavior. *Pharmacology Biochemistry Behavior* 15, 883-885.

Gorwood, P., 2012. Anxiety disorders and circadian rhythms. *Medicographia* 34, 289-294.

Graf, W.D., Oleinik, O.E., Glauser, T.A, Maertens, P., Eder, D.N., Pippenger, C.E., 1998. Altered antioxidant activities in children with a serious side adverse experience related to valproic acid therapy. *Neuropediatrics* 29, 195–201.

Graham, J.S., Beran, R.G., Kennaway, D.C., Mc Loughney, J., Matthews, C.D., 1995. Melatonin response in active epilepsy. *Epilepsia* 36,75–78.

Guardiola-Lemaître, B., Lenègre, A., Porsolt, R.D., 1992. Combined effects of diazepam and melatonin in two tests for anxiolytic activity in the mouse. *Pharmacology Biochemistry Behavior* 41, 405-408.

Gupta, M., Gupta, Y.K., Agarwal, S., Aneja, S., Kalaivani, M., Kohli, K., 2004. Effects of add-on melatonin administration on antioxidant enzymes in children with epilepsy taking

carbamazepine monotherapy: a randomized, double-blind, placebo-controlled trial. *Epilepsia* 45, 1636-1639.

Gupta, M., Aneja, S., Kohli, K., 2005. Add-on melatonin improves sleep behavior in children with epilepsy: randomized, double-blind, placebo-controlled trial. *Journal of Children Neurology* 20, 112-115.

Gupta, M., Kohli, K., Gupta, Y.K., 2006. Modulation of serum concentrations of melatonin by carbamazepine and valproate. *Indian Journal of Physiology and Pharmacology* 50, 79-82.

Jan, J.F., Tai, J., Hahn, G., Rothstein, R.R., 2001. Melatonin therapy for circadian rhythm sleep disorders in children with multiple disabilities: what have we learned in the last decade? *Developmental Medicine & Child Neurology* 46, 776–782.

Janjoppi, L., Silva de Lacerda, A.F., Scorza, F.A., Amado, D., Cavaleiro, E.A., Arida, R.M., 2006. Influence of pinealectomy on the amygdala kindling development in rats. *Neuroscience Letter* 392, 150-153.

Jones C, Huyton M, Hindley D., 2005. Melatonin and epilepsy. *Archives in Diseases in Childhood*. 90, 1203.

Juszcak, M., Drobnik, J., Guzek, J.W., Schwarzbarg, H., 1996. Effect of pinealectomy and melatonin on vasopressin-potentiated passive avoidance in rats. *Journal of Physiology and Pharmacology* 47, 621–627.

Kanner, A.M., Nieto, J.C., 1999. Depressive disorders in epilepsy. *Neurology* 53, S26-32.

Karakaş, A., Coskun, H., Kaya, A., Küçük, A., Gündüz B., 2011. The effects of the intraamygdalar melatonin injections on the anxiety like behavior and the spatial memory performance in male Wistar rats. *Behavioural Brain Research* 222, 141–150.

Kasper, S., Hajak, G., 2013. The efficacy of agomelatine in previously-treated depressed patients. *European Neuropsychopharmacology* 23, 814–821.

Kilic, E., Kilic, U., Bacigaluppi, M., Guo, Z., Abdallah, N.B., Wolfer, D.P., Reiter, R.J., Hermann, D.M., Bassetti, C.L., 2008. Delayed melatonin administration promotes neuronal survival, neurogenesis and motor recovery, and attenuates hyperactivity and anxiety after mild focal cerebral ischemia in mice. *Journal of Pineal Research* 45, 142-148.

Khezri, M.B., Oladi, M.R., Atlasbaf, A., 2013. Effect of melatonin and gabapentin on anxiety and pain associated with retrobulbar eye block for cataract surgery: a randomized double-blind study. *Indian Journal of Pharmacology* 45, 581-586.

Koh, S., Magid, R., Chung, H., Stine, C.D., Wilson, D.N., 2007. Deepressive behavior and selective down-regulation of serotonin receptor expression after early-life seizures: reversal by environmental enrichment. *Epilepsy & Behavior* 10, 26-31.

Konakchieva, R., Mitev, Y., Almeida, O., Patchev, V.K., 1998. Chronic melatonin treatment counteracts glucocorticoid-induced dysregulation of the hypothalamic-pituitary-adrenal axis in the rat. *Neuroendocrinology* 67, 171-180.

Kopp, C., Vogel, E., Rettori, M.C., Delagrang, P., Guardiola-Lemaître, B., Misslin, R., 1999. Effects of melatonin on neophobic responses in different strains of mice. *Pharmacology Biochemistry and Behavior* 63, 521-526.

Kopp, C., Vogel, E., Rettori, M., Delagrang, P., Misslin, R., 2000. Anxiolytic-like properties of melatonin receptor agonists in mice: involvement of MT1 and/or MT2 receptors in the regulation of emotional responsiveness. *Neuropharmacology* 39, 1865-1871.

- Kumar, A., Kaur, G., Rinwa, P., 2014. Buspirone along with melatonin attenuates oxidative damage and anxiety-like behavior in a mouse model of immobilization stress. *Chinese Journal of Natural Medicine* 12, 582-589.
- Lacoste, B., Angeloni, D., Dominguez-Lopez, S., Calderoni, S., Mauro, A., Fraschini, F., Descarries, L., Gobbi, G., 2015. Anatomical and cellular localization of melatonin MT1 and MT2 receptors in the adult rat brain. *Journal of Pineal Research* 58, 397-417.
- Lapin, I.P., Mirzaev, S.M., Ryzov, I.V., Oxenkrug, G.F., 1998. Anticonvulsant activity of melatonin against seizures induced by quinolinate, kainate, glutamate, NMDA, and pentylenetetrazole in mice. *Journal of Pineal Research* 24, 215-218.
- Laryea, G., Schütz, G., Muglia, L.J., 2013. Disrupting hypothalamic glucocorticoid receptors causes HPA axis hyperactivity and excess adiposity. *Molecular Endocrinology* 27, 1655-1665.
- Laudon, M., Urade, Y., Huang, Z., Neu-P., 2008. A novel melatonin agonist: effects on sleep and EEG power spectra in rats. *Sleep* 31 Suppl A34–35.
- Li, G., Li, P., Yang, X.L., 2001. Melatonin modulates gamma-aminobutyric acid(A) receptor-mediated currents on isolated carp retinal neurons. *Neuroscience Letters* 301, 49-53.
- Liu, C., Weaver, D.R., Jin, X., Shearman, L.P., Pieschl, R.L., Gribkoff, V.K., Reppert, S.M., 1997. Molecular dissection of two distinct actions of melatonin on the suprachiasmatic circadian clock. *Neuron*. 19, 91-102.
- Loiseau, F., Le Bihan, C., Hamon, M., Thiébot, M.H., 2006. Effects of melatonin and agomelatine in anxiety-related procedures in rats: interaction with diazepam. *European Neuropsychopharmacology* 16, 417-428.

Löscher, W., Fiedler, M., 1996. The role of technical, biological and pharmacological factors in the laboratory evaluation of anticonvulsant drugs. VI. Seasonal influences on maximal electroshock and pentylenetetrazol seizure thresholds. *Epilepsy Research* 25, 3-10.

Ma, J., Leung, L.S., 2004. Schizophrenia-like behavioral changes after partial hippocampal kindling. *Brain Research* 997, 111-118.

Maes, M., Bosmans, E., Meltzer, H.Y., Scharpé Suy, E., 1993. Interleukin-1 beta: a putative mediator of HPA axis hyperactivity in major depression? *The American Journal of Psychiatry* 150, 1189-1193.

Mairesse, J., Silletti, V., Laloux, C., Zuena, A.R., Giovine, A., Consolazione, M., van Camp, G., Malagodi, M., Gaetani, S., Ciani, S., Catalani, A., Mennuni, G., Mazzetta, A., van Reeth, O., Gabriel, C., Mocaër, E., Nicoletti, F., Morley-Fletcher, S., Maccari, S., 2013. Chronic agomelatine treatment corrects the abnormalities in the circadian rhythm of motor activity and sleep/wake cycle induced by prenatal restraint stress in adult rats. *International Journal of Neuropsychopharmacology* 16, 323-338.

Manev, H., Uz, T., Kharlamov, A., Joo, J.Y., 1996. Increased brain damage after stroke or excitotoxic seizures in melatonin-deficient rats. *FASEB journal* 10, 1546-1551.

Manni, R., Terzaghi, M., 2010. Comorbidity between epilepsy and sleep disorders. *Epilepsy Research* 90, 171-177.

Mantovani, M., Pértile, R., Calixto, J.B., Santos, A.R., Rodrigues, A.L., 2003. Melatonin exerts an antidepressant-like effect in the tail suspension test in mice: evidence for involvement of N-methyl-D-aspartate receptors and the L-arginine-nitric oxide pathway. *Neuroscience Letters* 343, 1-4.

Mazarati, A., Siddarth, P., Baldwin, R.A., Don, S., Rochelle, C., Raman, S., 2008. Depression after status epilepticus: behavioural and biochemical deficits and effects of fluoxetine. *Brain* 131, 2071-2083.

Mazarati, A.M., Shin, D., Kwon, Y.S., Bragin, A., Pineda, E., Tio, D., Taylor, A.N., Sankar, R., 2009. Elevated plasma corticosterone level and depressive behavior in experimental temporal lobe epilepsy. *Neurobiology of Diseases* 34, 457-461.

McIntyre, I.M., Norman, T.R., Marriott, P.F., Burrows, G.D., 1987. The pineal hormone melatonin in panic disorder. *Journal of Affective Disorders* 12, 203-206.

Meltzer, H., 1989. Serotonergic dysfunction in depression. *British Journal of Psychiatry* Suppl. 8, 25-31.

Mevissen, M., Ebert, U., 1998. Anticonvulsant effects of melatonin in amygdala-kindled rats. *Neuroscience Letters* 257, 13-16.

Micale, V., Arezzi, A., Rampello, L., Drago, F., 2006. Melatonin affects the immobility time of rats in the forced swim test: the role of serotonin neurotransmission. *European Neuropsychopharmacology* 16, 538-545.

Millan, M.J., Brocco, M., Gobert, A., Dekeyne, A., 2005. Anxiolytic properties of agomelatine, an antidepressant with melatonergic and serotonergic properties: role of 5-HT_{2C} receptor blockade. *Psychopharmacology (Berlin)* 177, 448-458.

Mittal, V.A., Karlsgodt, K., Zinberg, J., Cannon, T.D., Bearden, C.E., 2010. Identification and treatment of a pineal region tumor in an adolescent with prodromal psychotic symptoms. *American Journal of Psychiatry* 167, 1033-1037.

Moezi, L., Shafaroodi, H., Hojati, A., Dehpour, A.R., 2011. The interaction of melatonin and agmatine on pentylenetetrazole-induced seizure threshold in mice. *Epilepsy & Behavior* 22: 200-206.

Morley-Fletcher, S., Mairesse, J., Soumier, A., Banasr, M., Fagioli, F., Gabriel, C., Mocaer, E., Daszuta, A., McEwen, B., Nicoletti, F., Maccari, S., 2011. Chronic agomelatine treatment corrects behavioral, cellular, and biochemical abnormalities induced by prenatal stress in rats. *Psychopharmacology (Berlin)* 217, 301-313.

Nair, N.P., Hariharasubramanian, N., Pilapil, C., 1984. Circadian rhythm of plasma melatonin in endogenous depression. *Progress in Neuropsychopharmacology and Biological Psychiatry* 8, 715-718.

Naranjo-Rodriguez, E.B., Osornio, A.O., Hernandez-Avitia, E., Mendoza-Fernández, V., Escobar, A., 2000. Anxiolytic-like actions of melatonin, 5-metoxytryptophol, 5-hydroxytryptophol and benzodiazepines on a conflict procedure. *Progress in Neuropsychopharmacology and Biological Psychiatry* 24, 117-129.

Nava, F., Carta, G., 2001. Melatonin reduces anxiety induced by lipopolysaccharide in the rat. *Neuroscience Letters* 307, 57-60.

Nenchovska, Zl., Kortenska, L., Stefanova, M., Alova, L., Atanasova, M., Tchekalarova, J., 2014. Effects of pinealectomy on anxiety and depressive-like behavior in Wistar rats. *Comptes Rendus L'Academie Bulgarie des Sciences* 67, 1691-1700.

Niketic, V., Ristic, S., Saicic, Z.S., Spasic, M., Buzadzic, B., Stojkovic, M., 1995. Activities of antioxidant enzymes and formation of the glutathione adduct of hemoglobin (Hb ASSG) in epileptic patients with long-term antiepileptic therapy. *Farmaco*. 50, 811–813.

Niles, L.P., Pickering, D.S., Arciszewski, M.A., 1987. Effects of chronic melatonin administration on GABA and diazepam binding in rat brain. *Journal of Neural Transmission* 70, 117-124.

Niles, L.P., Peace, C.H., 1990. Allosteric modulation of t-[35S]butylbicyclophosphorothionate binding in rat brain by melatonin. *Brain Resarch Bulletin* 24, 635-638.

Ochoa-Sanchez, R., Rainer, Q., Comai, S., 2012. Anxiolytic effects of the melatonin MT(2) receptor partial agonist UCM765: comparison with melatonin and diazepam. *Progress in Neuropsychopharmacology and Biological Psychiatry* 39, 318-325.

Ott, D., Siddarth, P., Gurbani, S., Koh, S., Tournay, A., Shields, W.D., et al. 2003. Behavioral disorders in pediatric epilepsy: unmet psychiatric need. *Epilepsia* 44, 591–597.

O'Toole, K.K., Hooper, A., Wakefield, S., Maguire, J., 2014. Seizure-induced disinhibition of the HPA axis increases seizure susceptibility. *Epilepsy Research* 108, 29-43.

Overstreet, D.H., Pucilowski, O., Retton, M.C., Delagrange, P., Guardiola-Lemaitre, B., 1998. Effects of melatonin receptor ligands on swim test immobility. *Neuroreport* 9, 249-253.

Oxenkrug, G.F., Bachurin, S.O., Prakhie, IV, Zefirov, N.S., 2010. Quinone reductase 2 and antidepressant effect of melatonin derivatives. *Annals of New York Academy of Sciences* 1199, 121-124.

Païzanis, E., Renoir, T., Lelievre, V., Saurini, F., Melfort, M., Gabriel, C., Barden, N., Mocaër, E., Hamon, M., Lanfumey, L., 2010. Behavioural and neuroplastic effects of the new-generation antidepressant agomelatine compared to fluoxetine in glucocorticoid receptor-impaired mice. *International Journal of Neuropsychopharmacology* 13, 759-774.

Papp, M., Gruca, P., Boyer, P.A., Mocaer, E., 2003. Effect of Agomelatine in the Chronic Mild Stress Model of Depression in the Rat. *Neuropsychopharmacology* 28, 694–703.

Papp, M., Litwa, E., Gruca, P., Mocaer, E., 2006. Anxiolytic-like activity of agomelatine and melatonin in three animal models of anxiety. *Behavioral Pharmacology* 17, 9-18.

Paradiso, S., Hermann, B.P., Blumer, D., Davies, K., Robinson, R.G., 2001. Impact of depressed mood on neuropsychological status in temporal lobe epilepsy. *Journal of Neurology and Neurosurgery Psychiatry* 70, 180-185.

Pariante, C.M., Lightman, S.L., 2008. The HPA axis in major depression: classical theories and new developments. *Trends in Neuroscience* 31, 464-468.

Peled, N., Shorer, Z., Peled, E., Pillar, G., 2001. Melatonin effect on seizures in children with severe neurologic deficit disorders. *Epilepsia* 42, 1208-1210.

Petkova, Zl., Tchekalarova, J., Pechlivanova, D., Moyanova, Sl., Kortenska, L., Mitreva, R., Popov, D., Markova, P., Lozanov, V., Atanasova, D., Lazarov, N., Stoynev, Al., 2014. Treatment with melatonin after status epilepticus attenuates seizure activity and neuronal damage but does not prevent the disturbance in diurnal rhythms and behavioral alterations in spontaneously hypertensive rats in kainate model of temporal lobe epilepsy, *Epilepsy & Behavior* 31, 198-208.

Pévet, P., Bothorel, B., Slotten, H., Saboureau, M., 2002. The chronobiotic properties of melatonin. *Cell Tissue Research* 309, 183-191.

Pévet, P., Challet, E., 2011. Melatonin: both master clock output and internal time-giver in the circadian clocks network. *Journal of Physiology (Paris)* 105, 170-182.

Pierrefiche, G., Zerbib, R., Laborit, H., 1993. Anxiolytic activity of melatonin in mice: involvement of benzodiazepine receptors. *Research Communication and Chem Pathology and Pharmacology* 82, 131-142.

Pineda, E., Shin, D., Sankar, R., Mazarati, A.M., 2010. Comorbidity between epilepsy and depression: experimental evidence for the involvement of serotonergic, glucocorticoid, and neuroinflammatory mechanisms. *Epilepsia* 51 Suppl 3, 110-114.

Pineda, E., Jentsch, J.D., Shin, D., Griesbach, G., Sankar, R., Mazarati, A., 2014. Behavioral impairments in rats with chronic epilepsy suggest comorbidity between epilepsy and attention deficit/hyperactivity disorder. *Epilepsy & Behavior* 31, 267-275.

Prakhie, IV., Oxenkrug, G.F., 1998. The effect of nifedipine, Ca(2+) antagonist, on activity of MAO inhibitors, N-acetylserotonin and melatonin in the mouse tail suspension test. *International Journal of Neuropsychopharmacology* 1, 35-40.

Quigg, M., 2000. Circadian rhythms: interactions with seizures and epilepsy. *Epilepsy Research* 42, 43-55.

Raedt, R., Van Dycke, A., Van Melkebeke, D., De Smedt, T., Claeys, P., Wyckhuys, T., Vonck, K., Wadman, W., Boon, P., 2009. Seizures in the intrahippocampal kainic acid epilepsy model: characterization using long-term video-EEG monitoring in the rat. *Acta Neurologica Scandinavica* 119, 293-303.

Raghavendra, V., Kaur, G., Kulkarni, S.K., 2000. Anti-depressant action of melatonin in chronic forced swimming-induced behavioral despair in mice, role of peripheral benzodiazepine receptor modulation. *European Neuropsychopharmacology* 10, 473-481.

Ramírez-Rodríguez, G., Vega-Rivera, N.M., Oikawa-Sala, J., Gómez-Sánchez, A., Ortiz-López, L., Estrada-Camarena, E., 2014. Melatonin synergizes with citalopram to induce

antidepressant-like behavior and to promote hippocampal neurogenesis in adult mice. *Journal of Pineal Research* 56, 450-461.

Ray, M., Mediratta, P.K., Reeta, Kh., Mahajan, P., Sharma, K.K., 2004. Receptor mechanisms involved in the anticonvulsant effect of melatonin in maximal electroshock seizures. *Methods and Findings in Experimental and Clinical Pharmacology* 26, 177-181.

Richerson, G.B., Buchanan, G.F., 2011. The serotonin axis: Shared mechanisms in seizures, depression, and SUDEP. *Epilepsia* 52 Suppl 1, 28-38.

Rosenstein, R.E., Cardinali, D.P., 1986. Melatonin increases in vivo GABA accumulation in rat hypothalamus, cerebellum, cerebral cortex and pineal gland. *Brain Research* 398, 403–406.

Rosenstein, R.E., Estevez, A.G., Cardinali, D.P., 1989. Time-dependent effect of melatonin on glutamic acid decarboxylase activity and $^{36}\text{Cl}^-$ influx in rat hypothalamus. *Journal of Neuroendocrinology* 1, 443–447.

Salva, M.A., Hartley, S., 2012. Mood disorder, circadian rhythms, melatonin and melatonin agonists. *Journal of Central Nervous System Disorders* 4, 15-26.

Sandyk, R., Tsagas, N., Anninos, P.A., 1992. Melatonin as a proconvulsive hormone in humans. *International Journal of Neuroscience* 63, 125-135.

Sansone, R.A., Sansone, L.A., 2011. Agomelatine: a novel antidepressant. *Innovative Clinical Neuroscience* 8, 10-14.

Serfaty, M.A., Osborne, D., Buszewicz, M.J., Blizard, R., Raven, P.W., 2010. A randomized double-blind placebo-controlled trial of treatment as usual plus exogenous slow-release

melatonin (6 mg) or placebo for sleep disturbance and depressed mood. *International Clinical Psychopharmacology* 25, 132-142.

Silva de Lacerda, A.F., Janjoppi, L., Scorza, F.A., Lima, E., Amado, D., Cavaleiro, E.A., Arida, R.M., 2007. Physical exercise program reverts the effects of pinealectomy on the amygdala kindling development. *Brain Research Bulletin* 74, 216-220.

Shaji, A.V., Kulkarni, S.K., 1998. Central nervous system depressant activities of melatonin in rats and mice. *Indian Journal of Experimental Biology* 36, 257-263.

Shouse, M.N., da Silva, M., Sammaritano, A.M., 1996. Circadian Rhythm, Sleep, and Epilepsy. *Journal of Clinical Neurophysiology* 13, 32-50.

Solmaz, I., Gürkanlar, D., Gökçil, Z., Göksoy, C., Ozkan, M., Erdoğan, E., 2009. Antiepileptic activity of melatonin in guinea pigs with pentylenetetrazol-induced seizures. *Neurological Research* 31, 989-995.

Staba, R.J., Wilson, C.L., Bragin, A., Fried, I., Engel, J. Jr., 2002. Sleep states differentiate single neuron activity recorded from human epileptic hippocampus, entorhinal cortex, and subiculum. *Journal of Neuroscience* 22, 5694-5704.

Stahl, S.M., 2014. Mechanism of action of agomelatine: a novel antidepressant exploiting synergy between monoaminergic and melatonergic properties. *CNS Spectrums* 19, 207-212.

Stein, D.J., Ahokas, A.A., de Bodinat, C., 2008. Efficacy of agomelatine in generalized anxiety disorder: a randomized, double-blind, placebo-controlled study. *Journal of Clinical Psychopharmacology* 28, 561-566.

Stewart, L.S., 2001. Endogenous melatonin and epileptogenesis: facts and hypothesis. *International Journal of Neuroscience* 107, 77-85.

Stewart, L.S., Leung, L.S., 2003. Temporal lobe seizures alter the amplitude and timing of rat behavioral rhythms. *Epilepsy & Behavior* 4, 153-160.

Sumay, I.C., Masana, M.I., Dubocovich, M.L., 2005. The antidepressant-like effect of the melatonin receptor ligand luzindole in mice during forced swimming requires expression of MT₂ but not MT₁ melatonin receptors. *Journal of Pineal Research* 39, 170-177.

Tchekalarova, J., Pechlivanova, D., Itzev, D., Lazarov, N., Markova, P., Stoynev, A., 2010. Diurnal rhythms of spontaneous recurrent seizures and behavioral alterations of Wistar and spontaneously hypertensive rats in the kainate model of epilepsy. *Epilepsy & Behavior* 17, 23-32.

Tchekalarova, J., Petkova, Z., Pechlivanova, D., Moyanova, Sl., Kortenska, L., Mitreva, R., Lozanov, V., Atanasova, D., Lazarov, N., Stoynev, Al., 2013. Prophylactic treatment with melatonin after status epilepticus: Effects on epileptogenesis, neuronal damage and behavioral changes in kainate model of temporal lobe epilepsy. *Epilepsy & Behavior* 27, 174-187.

Tchekalarova, J., 2014. Finding a model for comorbidity between attention deficit hyperactivity disorder and epilepsy. *Molecular and Cellular Epilepsy* 1, 92-95.

Tian, S., Laudon, M., Han, L., Gao, J., Huang, J., Yang, Y., Deng, H., 2010. Antidepressant- and anxiolytic effects of the novel melatonin agonist Neu-P11 in rodent models. *Acta Pharmacologica Sinica* 31, 775–783.

Turek, F.W., 2007. From circadian rhythms to clock genes in depression. *International Clinical Psychopharmacology* 22 Suppl 2, S1-8.

Upton, N., Stean, T., Middlemiss, D., Blackburn, T., Kennett, G., 1998. Studies on the role of 5-HT_{2C} and 5-HT_{2B} receptors in regulating generalised seizure threshold in rodents. *European Journal of Pharmacology* 359, 33-40.

Vimala, P.V., Bhutada, P.S., Patel, F.R., 2014. Therapeutic potential of agomelatine in epilepsy and epileptic complications. *Medicinal Hypotheses* 82, 105-110.

Wade, A.G., Ford, I., Crawford, G., McConnachie, A., Nir, T., Laudon, M., Zisapel, N., 2010. Nightly treatment of primary insomnia with prolonged release melatonin for 6 months: a randomized placebo controlled trial on age and endogenous melatonin as predictors of efficacy and safety. *BMC Medicine* 8, 51.

Wang, F., Li, J., Wu, C., Yang, J., Xu, F., Zhao, Q., 2003. The GABA(A) receptor mediates the hypnotic activity of melatonin in rats. *Pharmacology Biochemistry and Behavior* 74, 573-578.

Weil, Z.M., Hotchkiss, A.K., Gatien, M.L., Pieke-Dahl, S., Nelson, R.J., 2006. Melatonin receptor (MT1) knockout mice display depression-like behaviors and deficits in sensorimotor gating. *Brain Research Bulletin* 68, 425-429.

Weiss, G., Lucero, K., Fernandez, M., Karnaze, D., Castillo, N., 1993. The effect of adrenalectomy on the circadian variation in the rate of kindled seizure development. *Brain Research* 612, 354-356.

Weidenfeld, Y., Schmidt, U., Nir, I., 1993. The effect of exogenous melatonin on the hypothalamic-pituitary-adrenal axis in intact and pinealectomized rats under basal and stressed conditions. *Journal of Pineal Research* 14, 60-66.

Witt-Enderby, P.A., Bennett, J., Jarzynka, M.J., Firestine, S., Melan, M.A., 2003. Melatonin receptors and their regulation: biochemical and structural mechanisms. *Life Science* 72, 2183-2198.

Wu, F.S., Yang, Y.C., Tsai, J.J., 1999. Melatonin potentiates the GABA(A) receptor-mediated current in cultured chick spinal cord neurons. *Neurosci Letters* 260, 177-180.

- Yahyavi-Firouz-Abadi, N., Tahsili-Fahadan, P., Riazi, K., Ghahremani, M.H., Dehpour, A.R., 2007. Melatonin enhances the anticonvulsant and proconvulsant effects of morphine in mice: role for nitric oxide signaling pathway. *Epilepsy Research* 75, 138-144.
- Yalın, O., Arman, F., Erdoğan, F., Kula, M., 2006. A comparison of the circadian rhythms and the levels of melatonin in patients with diurnal and nocturnal complex partial seizures. *Epilepsy & Behavior* 8, 542-546.
- Yehuda, S., Mostofsky, D.I., 1993. Circadian effects of beta-endorphin, melatonin, DSIP, and amphetamine on pentylenetetrazol-induced seizures. *Peptides* 14, 203-205.
- Yildirim, M., Aydin-Abidin, S., Abidin, I., Akca, M., Canpolat, S., Cansu, A., 2013. Evaluation of the role of chronic daily melatonin administration and pinealectomy on penicillin-induced focal epileptiform activity and spectral analysis of ECoG in rats: an in vivo electrophysiological study. *Neurochemistry Research* 38, 1672-1685.
- Yonei Y., Hattori, A., Tsutsui, K., Okawa, M., Ishizuka, B., 2010. Effects of Melatonin: Basics Studies and Clinical Applications. *Anti-Aging Medicine. Japanese Society of Anti-Aging Medicine* 7, 85-91.
- Yousaf, F., Seet, E., Venkatraghavan, L., Abrishami, A., Chung, F., 2010. Efficacy and safety of melatonin as an anxiolytic and analgesic in the perioperative period: a qualitative systematic review of randomized trials. *Anesthesiology* 113, 968-976.
- Yüksel, A., Cengiz, M., Seven, M., Ulutin, T., 2000. Erythrocyte glutathione, glutathione peroxidase, superoxide dismutase and serum lipid peroxidation in epileptic children with valproate and carbamazepine monotherapy. *Journal of Basic Clinical Physiology and Pharmacology* 11, 73-81.

Zahra, E.M., Ibtissam, L., Abdelhalim, M., Aboubakr, E.H., Ouichou, A., 2012. Effects of Melatonin in Male and Female Wistar Rats: A Possible Implication of Sex Hormones. Neuroscience & Medicine 3, ID:19649,12 pages.